Converse KAM Theory for monotone positive symplectomorphisms

Àlex Haro

Departament de Matemàtica Aplicada i Anàlisi, Facultat de Matemàtiques, Universitat de Barcelona. Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain. E-mail: haro@cerber.mat.ub.es

Abstract. We apply variational methods to Converse KAM theory. These are useful for symplectomorphisms in the annulus that satisfy weaker hypotheses than those usually required. For instance, we do not need the existence of a global Lagrangian generating function. We obtain the variational principles from the primitive function of our symplectomorphism. They are introduced not only for the orbits of a symplectomorphism, but also for the so-called invariant Lagrangian graphs. Among the non-degenerate i.L.g. we focus on the minimizing ones. Applications are also described for a broad class of examples.

AMS classification scheme numbers: 58F05

1. Introduction

A fundamental question in symplectic-Hamiltonian dynamics is which parts of the phase space contain invariant tori and which do not. Often one works on the *d-annulus* $\mathbb{A}^d$, that is, the cotangent bundle $T^*\mathbb{T}^d \simeq \mathbb{T}^d \times \mathbb{R}^d = \mathbb{A}^d$ of the d-torus $\mathbb{T}^d = \mathbb{R}^d/\mathbb{Z}^d$, or in its universal covering space $\mathbb{R}^d \times \mathbb{R}^d$.

- On the one hand, *KAM theory* (from Kolmogorov [18], Arnold [1] and Moser [26]) has enabled us to obtain *many* invariant tori of dimension d for the exact symplectomorphisms which are close enough to completely integrable ones. Some non-degeneracy conditions are needed. In fact, this theory proves that the measure of the complement of the invariant tori in any bounded region is arbitrarily small when the size of the perturbation is reduced. The dynamics on these tori are conjugated to ergodic translations on $\mathbb{T}^d$ satisfying Diophantine conditions. Herman has proved that these tori need to be Lagrangian [15].
- On the other hand, the Birkhoff's theory [8] concerning the invariant rotational curves for area preserving maps in the annulus $\mathbb{A} = \mathbb{T} \times \mathbb{R}$ is the first step towards a global (non-perturbative) study on the existence of such curves. This theory gives several Lipschitzian inequalities for the invariant curves and first asserts that they must be graphs (under some non-degeneracy conditions). These theorems also appear in the studies of Herman [14] and Mather [24]. Herman improved

and generalized these results to higher dimensions in the course of several papers [15, 16]. We shall refer to this theory as *BHM theory*. Bialy and Polterovich [7] have a high-dimensional version of the 'second' Birkhoff's theorem for optical Hamiltonian diffeomorphisms.

- Finally, it would be useful to determine the conditions under which there are no invariant tori through a given point or region in phase space. The development of such criteria is known as *Converse KAM theory*. After the earlier work by Aubry and Mather, non-existence criteria were found for area preserving twist maps [20] and later for symplectic twist maps [21]. A summary of these results appears in [9].

In order to study the existence of *invariant rotational tori* for symplectomorphisms defined on $\mathbb{T}^d \times \mathbb{R}^d$ (that is, invariant tori which are homotopic to the zero-section), here we apply variational methods, attempting to ensure that they are more effective in higher dimensions and under weaker hypotheses than those usually required (say, that our symplectomorphism satisfies a strong condition of positivity). Although we still need some positivity conditions and we are only able to work with those tori that are graphs, it might be possible that our results work in more general cases by using suitable coordinates, thanks to the local character of our variational principles. As a summary, we obtain that:

> *the orbits on a monotone positive invariant Lagrangian horizontal torus are minimizing.*

The definition of monotone positivity was given by Herman [15, 16], although he referred to symplectic maps rather than invariant tori, and the result is similar to that obtained by MacKay, Meiss and Stark [21]. We obtain the following *non-existence criterion* of invariant tori (cf. [21]):

> *If the orbit by z yields on a monotone positive region and it has a non-minimizing segment, then it does not lie on any invariant Lagrangian horizontal torus included within such a region.*

Notice that we do not require global conditions for our symplectomorphism because we can test whether a segment of orbit is a local minimum of the corresponding action. Hence, the existence of a global generating function, which involves global conditions for our symplectomorphism, is not necessary.

We have applied this to a broad class of examples, given by the family of generalized standard-like maps [19]. Some examples that appear in this paper are examples of non-twist symplectic maps, but we can study the region in phase space where the positivity conditions are satisfied.

2. Definitions

2.1. The symplectic group

We can write any matrix M of $M_{2d}(\mathbb{R})$ by $d\times d$ blocks:

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

Let J be the symplectic matrix

$$J = \begin{pmatrix} 0 & I_d \\ -I_d & 0 \end{pmatrix},$$

where I_d is the $d \times d$-identity matrix. $\mathrm{Sp}(2d)$ denotes the symplectic group of $\mathbb{R}^{2d}$ (subgroup of $\mathrm{GL}(2d,\mathbb{R})$), that is to say, the set of matrices $M \in M_{2d}(\mathbb{R})$ such that $M^\top JM = J$ (where $^\top$ means the transpose of a matrix). Hence:

$$M \in \mathrm{Sp}(2d) \Leftrightarrow M^\top JM = J \Leftrightarrow A^\top C = C^\top A, B^\top D = D^\top B, A^\top D - C^\top B = I_d,$$
$$\Updownarrow$$
$$M^\top \in \mathrm{Sp(2d)} \Leftrightarrow MJM^\top = J \Leftrightarrow AB^\top = BA^\top, CD^\top = DC^\top, AD^\top - BC^\top = I_d.$$

2.2. The primitive function of a symplectomorphism

We consider $\mathbb{R}^{2d} = \mathbb{R}^d \times \mathbb{R}^d$ endowed with the *position-momentum* coordinates

$$z = (x, y) = (x^1, \ldots, x^d, y^1, \ldots, y^d).$$

For the sake of simplicity, all the maps will be $\mathcal{C}^\infty$.

Definition: Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a diffeomorphism, that is, we write

$$\begin{cases} x' = f(x, y), \\ y' = g(x, y). \end{cases}$$

It is a *symplectomorphism* iff $\forall z \in \mathbb{R}^{2d}$, $\mathrm{D}F(z) \in \mathrm{Sp}(2d)$. In such a case, the *exactness equations* associated to F, which are the Pfaffian system

$$\begin{cases} \dfrac{\partial S}{\partial x}(x, y) = g(x, y)^\top \dfrac{\partial f}{\partial x}(x, y) - y^\top, \\ \\ \dfrac{\partial S}{\partial y}(x, y) = g(x, y)^\top \dfrac{\partial f}{\partial y}(x, y), \end{cases}$$

are solvable.

Definition: A solution $S : \mathbb{R}^{2d} \to \mathbb{R}$ of the exactness equations is referred to as a *primitive function* of F, and it is defined apart from additive constants. Anyway, we write $\mathrm{pf}(F) = S$.

Remarks:

i) Recall that the standard symplectic structure on $\mathbb{R}^{2d}$ is given by $\boldsymbol{\omega} = \mathrm{d}y \wedge \mathrm{d}x$, and it is exact: $\boldsymbol{\alpha} = y\,\mathrm{d}x$ is an action form. The exactness equations are the translation to coordinates of the condition $F^*\boldsymbol{\alpha} - \boldsymbol{\alpha} = \mathrm{d}S$. However, we shall not use this language in this paper.

ii) In the literature, the primitive function is often called generating function, although it does not generate the symplectomorphism! Because of this, we have followed the nomenclature used in [3]. Anyway, the primitive function and the different kinds of generating functions are related by certain p.d.e. [13].

iii) Symplectomorphisms are closed by composition and inversion. Given two symplectomorphisms $F, G : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$, with $\mathrm{pf}(F) = S$ and $\mathrm{pf}(G) = T$, we have the following formulae:

(i) $\mathrm{pf}(G \circ F) = S + T \circ F$,
(ii) $\mathrm{pf}(F^{-1}) = -S \circ F^{-1}$,
(iii) $\mathrm{pf}(G^{-1} \circ F \circ G) = S \circ G + T - T \circ G^{-1} \circ F \circ G$.

iv) Thanks to the topological properties of $\mathbb{R}^{2d}$, the primitive function of a symplectomorphism always exists. This is not necessary in more general contexts, and we require an exact symplectic structure (see, for instance [13]). ◁

Examples:

1) Any diffeomorphism of the form

$$\begin{pmatrix} x \\ y \end{pmatrix} \to \begin{pmatrix} \phi(x) \\ \mathrm{D}\phi(x)^{-\top} y \end{pmatrix},$$

where $\phi : \mathbb{R}^d \to \mathbb{R}^d$ is a diffeomorphism, is a symplectomorphism and has primitive function equal to zero ($^{-\top}$ means the transpose of the inverse of a matrix). This symplectomorphism is the *lift* of the diffeomorphism ϕ, and it is represented by $\hat{\phi}$. It is easy to see that if we compose on the left our initial symplectomorphism F with a lift $L = \hat{\phi}$ then we obtain another symplectomorphism $\bar{F} = L \circ F$ with the same primitive function.

2) A *(symplectic) vertical translation* $\tau = \tau_{\nabla l}$ is defined by means of a function $l : \mathbb{R}^d \to \mathbb{R}$ and it is given by

$$\begin{pmatrix} x \\ y \end{pmatrix} \to \begin{pmatrix} x \\ y + \nabla l(x) \end{pmatrix}.$$

It is a symplectomorphism and its primitive function is just l (it is a function which only depends on the x-variables). ◁

2.3. Monotonicity

Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism. We write

$$\mathrm{D}F(z) = \begin{pmatrix} A(z) & B(z) \\ C(z) & D(z) \end{pmatrix}.$$

Definition: We say that F is *monotone* iff $\forall z \in \mathbb{R}^{2d}$ $|B(z)| \neq 0$. In such a case, the matrices $B^{-1}(z)A(z)$ and $D(z)B^{-1}(z)$ are symmetric. Following [15, 16], we shall say that F is *monotone positive* iff one or both of the following conditions is verified:

$(+_a)$ $\forall z \in \mathbb{R}^{2d}$ $B^{-1}(z)A(z)$ is positive definite;

$(+_d)$ $\forall z \in \mathbb{R}^{2d}$ $D(z)B^{-1}(z)$ is positive definite.

We shall distinguish between the two types of monotone positivity by writing $(+_a)$ or $(+_d)$. We define monotone negativity in the same way. We can perform analogous definitions for monotone regions, monotone points, etc.

The symmetric matrix

$$T(z) = \frac{1}{2}(B(z) + B(z)^\top)$$

is the *torsion* of F at the point z. If it is positive definite for all the points, we shall say that our diffeomorphism *has positive torsion*. If the torsion is uniformly positive definite, we shall say that F is a *twist map* and, as Avez proved in [6] (see also [21]), the map

$$\begin{pmatrix} x \\ y \end{pmatrix} \to \begin{pmatrix} x \\ f(x, y) \end{pmatrix}$$

is a diffeomorphism on $\mathbb{R}^{2d}$.

Remark: Notice that symplectic matrices exist which are monotone positive but which do not have positive torsion, and others that have positive torsion and are not monotone positive. Simple examples, for $d = 1$, are given by

$$\begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}.$$

◁

2.4. Generating functions

It is well known that, sometimes, a symplectomorphism is given by a (Lagrangian) generating function. We need monotonicity. In fact, we need a stronger condition.

Definition: Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a monotone symplectomorphism, and $\mathrm{pf}(F) = S$. We say that it is *strongly monotone* iff

$$\forall x \in \mathbb{R}^d, f(x, \cdot) : \mathbb{R}^d \to \mathbb{R}^d \text{ is a diffeomorphism.}$$

Symplectic twist maps are examples of strongly monotone symplectomorphisms [6, 21].

In such a case, let $\phi(x, \cdot)$ be the inverse of $f(x, \cdot)$, that is, $\forall x, x' \in \mathbb{R}^d$ $x' = f(x, \phi(x, x'))$. We define the function $L : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ by

$$L(x, x') = S(x, \phi(x, x')).$$

Therefore, applying the exactness equations, we obtain the relations

$$\begin{cases} y = -\nabla_x L(x, x'), \\ y' = \nabla_{x'} L(x, x'), \end{cases}$$

which define implicitly our symplectomorphism. The function L is called a *(global) Lagrangian generating function* of F, and defines implicitly our symplectomorphism F.

Hence, the relationship between the Lagrangian generating function and the primitive function is given by

$$S\left(x, -\nabla_x L(x, x')\right) = L(x, x').$$

Moreover, the second derivatives of L are given by

$$\frac{\partial^2 L}{\partial x^2} = B^{-1}A, \ \frac{\partial^2 L}{\partial x' \partial x} = -B^{-1}, \ \frac{\partial^2 L}{\partial x'^2} = DB^{-1}.$$

Remarks:

i) Notice that these expressions appear in the definitions of monotonicity.

ii) If one asks for the inverse problem, that is, to obtain the generating function from the primitive function, one has to solve this p.d.e., and we need some additional information. ◁

An orbit $Z = (x_k, y_k)_{k \in \mathbb{Z}}$ is determined by the positions $(x_k)_{k \in \mathbb{Z}}$. Formally speaking, the orbits are in correspondence with the stationary bisequences $X = (x_k)_{k \in \mathbb{Z}}$ of the action

$$\boldsymbol{L}(X) = \sum_{k \in \mathbb{Z}} L(x_k, x_{k+1})$$

(see [5, 22]). Notice also that the value of this action along an orbit is $\sum_{k \in \mathbb{Z}} S(x_k, y_k)$.

3. Variational principles

Since the generating function of a symplectomorphism is not always well defined, or its computation is difficult, we shall perform variational principles without its aid. The final part of this paper is devoted to some examples of symplectomorphisms which are not defined by means of generating functions.

3.1. Chains and connecting orbits

Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism with $\mathrm{pf}(F) = S$. Given two x-points $\mathbf{x}_m, \mathbf{x}_n \in \mathbb{R}^d$, where $n > m+1$ are two indices, we want to search for the *connecting orbits* from $\mathbf{x}_m$ to $\mathbf{x}_n$, that is, the (m, n)-segments of $\mathbb{R}^{2d}$

$$(x_m, y_m), (x_{m+1}, y_{m+1}), \ldots (x_{n-1}, y_{n-1})$$

such that

- $x_m = \mathbf{x}_m$, $f(x_{n-1}, y_{n-1}) = \mathbf{x}_n$,
- $\forall i = m \div n-2$ $F(x_i, y_i) = (x_{i+1}, y_{i+1})$.

We shall consider the (m, n)-*orbital action*

$$\boldsymbol{S}_{m,n}(x_m, y_m, x_{m+1}, y_{m+1}, \ldots, x_{n-1}, y_{n-1}) = \sum_{i=m}^{n-1} S(x_i, y_i)$$

restricted to the (m, n)-segments satisfying

- $x_m = \mathbf{x}_m$, $f(x_{n-1}, y_{n-1}) = \mathbf{x}_n$,
- $\forall i = m \div n-2$ $f(x_i, y_i) = x_{i+1}$.

They are the *chains* from $\mathbf{x}_m$ to $\mathbf{x}_n$, and the corresponding set of chains, $\mathcal{C}_{\mathbf{x}_m,\mathbf{x}_n}$, is a $d(n-m-1)$-submanifold of $\mathbb{R}^{2d(n-m)}$, provided the rank of the matrix

$$\begin{pmatrix} B_m & -I & & & & & \\ A_{m+1} & B_{m+1} & -I & & & & \\ & \ddots & \ddots & \ddots & & & \\ & & A_{n-2} & B_{n-2} & -I & & \\ & & & & A_{n-1} & B_{n-1} \end{pmatrix}$$

is maximal ($= d(n-m)$) in all the chains. For instance, if F is monotone this *transversality condition* is satisfied. In general, if this condition is satisfied for a certain connecting orbit, we can say that it is *transversal*. An orbit is transversal iff all its segments are transversal.

Proposition 1 *Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism, with $\mathrm{pf}(F) = S$. Given two x-points $\mathbf{x}_m, \mathbf{x}_n \in \mathbb{R}^d$, suppose that the corresponding set of chains, $\mathcal{C}_{\mathbf{x}_m,\mathbf{x}_n}$, satisfies the transversality condition, and consider the orbital action $\boldsymbol{S}_{m,n}$ on it. Then:*

- *The connecting orbits are critical chains of $\boldsymbol{S}_{m,n}$.*
- *If F is monotone, the critical chains of $\boldsymbol{S}_{m,n}$ are connecting orbits of F.*

Proof: By the Lagrange multiplier method, we have to search for the critical points of the function

$$L(y_m, x_{m+1}, y_{m+1}, \ldots, x_{n-1}, y_{n-1}, \lambda_{m+1}, \ldots, \lambda_n) = \sum_{i=m}^{n-1} (S(x_i, y_i) + \lambda_{i+1}(x_{i+1} - f(x_i, y_i))),$$

where all the variables belong to $\mathbb{R}^d$, and we have taken $x_m = \mathbf{x}_m$ and $x_n = \mathbf{x}_n$. The equations that we have to solve are

$$0 = \frac{\partial L}{\partial x_i} = (g(x_i, y_i) - \lambda_{i+1})^\top \frac{\partial f}{\partial x}(x_i, y_i) + (\lambda_i - y_i)^\top \qquad (i = m{+}1 \div n{-}1),$$

$$0 = \frac{\partial L}{\partial y_i} = (g(x_i, y_i) - \lambda_{i+1})^\top \frac{\partial f}{\partial y}(x_i, y_i) \qquad (i = m \div n{-}1),$$

$$0 = \frac{\partial L}{\partial \lambda_i} = (x_i - f(x_{i-1}, y_{i-1}))^\top \qquad (i = m{+}1 \div n),$$

where we have used the exactness equations (2.2). Therefore:

- If $(x_i, y_i)_{i=m\div n-1}$ is a connecting orbit from $\mathbf{x}_m$ to $\mathbf{x}_n$, then it is a critical chain of the function $\boldsymbol{S}_{m,n}$ (take $\forall i = m{+}1 \div n{-}1$ $\lambda_i = y_i$ and $\lambda_n = g(x_{n-1}, y_{n-1})$).
- Suppose that $\left|\frac{\partial f}{\partial y}\right| \neq 0$. For a critical chain of $\boldsymbol{S}_{m,n}$ ($\forall i = m \div n{-}1$ $x_{i+1} = f(x_i, y_i)$) the second set of equations gives $\forall i = m \div n{-}1$ $\lambda_{i+1} = g(x_i, y_i)$ and, therefore, the first one gives $\forall i = m{+}1 \div n{-}1$ $\lambda_i = y_i$. Finally, we obtain $\forall i = m \div n{-}2$ $g(x_i, y_i) = \lambda_{i+1} = y_{i+1}$.

□

Remarks:

i) We can extend these ideas to any cotangent bundle of a manifold $\mathcal{M}$, $T^*\mathcal{M}$, where a canonical exact symplectic structure is defined [2]. Recall that the cotangent bundles are the *phase spaces* of classical mechanics, and the basic manifolds $\mathcal{M}$ are the *configuration spaces*. For further information about these variational principles of *discrete mechanics* see [13].

ii) From a physical point of view, if our symplectomorphism is the time-1 map of a certain time-periodic Hamiltonian (with period 1), an F-chain corresponds to an 'orbit' whose velocity is crudely changed every period (as in a maneuver). Hence, the fact to make the action extreme on the set of F-chains corresponds to the smoothing of the peaks. ◁

3.2. Extremal character of an orbit

We have seen that the connecting orbits from $\mathbf{x}_m$ to $\mathbf{x}_n$ are critical F-chains of a certain action. We need to consider their extremal character, i.e., the character of the second derivative $H_{m,n} = \mathrm{D}^2 \boldsymbol{S}_{m,n}$.

If F is monotone, or at least it is monotone in the region where the segment lives, then the function $\boldsymbol{S}_{m,n}$ can be locally written in variables $x_{m+1}, \ldots, x_{n-1}$, and we can compute the corresponding Hessian matrix. That is, the set of equations $\{f(x_i, y_i) = x_{i+1}\}_{i=m\div n-1}$ defines implicitly a set of functions $\{\eta_i = \eta_i(x, x')\}_{i=m\div n-1}$ such that $\forall i = m \div n{-}1$

$$f(x_i, \eta_i(x_i, x_{i+1})) = x_{i+1}.$$

Of course, these functions are defined on a neighborhood of our connecting orbit. Their derivatives are given from

$$0 = \frac{\partial f}{\partial x}(x_i, \eta_i(x_i, x_{i+1})) + \frac{\partial f}{\partial y}(x_i, \eta_i(x_i, x_{i+1}))\ \frac{\partial \eta_i}{\partial x}(x_i, x_{i+1})$$

and

$$I = \frac{\partial f}{\partial y}(x_i, \eta_i(x_i, x_{i+1}))\ \frac{\partial \eta_i}{\partial x'}(x_i, x_{i+1}).$$

Therefore, we have to compute the critical points of the function

$$\boldsymbol{S}_{m,n}(x_1, \ldots, x_{n-1}) = \sum_{i=m}^{n-1} S(x_i, \eta_i(x_i, x_{i+1})),$$

where we have taken $x_m = \mathbf{x}_m$ and $x_n = \mathbf{x}_n$. So then, $\forall i = m+1 \div n-1$

$$\begin{aligned}\frac{\partial \boldsymbol{S}_{m,n}}{\partial x_i} &= \frac{\partial S}{\partial x}(x_i, \eta_i) + \frac{\partial S}{\partial y}(x_i, \eta_i)\ \frac{\partial \eta_i}{\partial x}(x_i, x_{i+1}) + \frac{\partial S}{\partial y}(x_{i-1}, \eta_{i-1})\ \frac{\partial \eta_{i-1}}{\partial x'}(x_{i-1}, x_i)\\ &= g_i^\top\ \frac{\partial f}{\partial x}(x_i, \eta_i) - \eta_i^\top + g_i^\top\ \frac{\partial f}{\partial y}(x_i, \eta_i)\ \frac{\partial \eta_i}{\partial x} + g_{i-1}^\top\ \frac{\partial f}{\partial y}(x_{i-1}, \eta_{i-1})\ \frac{\partial \eta_{i-1}}{\partial x'}\\ &= g_{i-1}^\top - \eta_i^\top,\end{aligned}$$

where $g_i = g(x_i, \eta_i)$. As we can see, our connecting orbit makes the action extreme. We are going to compute the second derivatives. We have written $A_i = \frac{\partial f}{\partial x}(x_i, y_i)$, etc. Then, $\forall i = m+1 \div n-1$

$$\frac{\partial^2 \boldsymbol{S}_{m,n}}{\partial x_i^2} = \frac{\partial g}{\partial y}(x_{i-1}, y_{i-1})\ \frac{\partial \eta_{i-1}}{\partial x'}(x_{i-1}, x_i) - \frac{\partial \eta_i}{\partial x}(x_i, x_{i+1}) = D_{i-1}\ B_{i-1}^{-1} + B_i^{-1}\ A_i$$

and $\forall i = m+1 \div n-2$

$$\frac{\partial^2 \boldsymbol{S}_{m,n}}{\partial x_{i+1} \partial x_i} = -\frac{\partial \eta_i}{\partial x'}(x_i, x_{i+1}) = -B_i^{-1}\ .$$

In summary:

Proposition 2 *The Hessian matrix associated to a monotone (m,n)-segment of orbit is given by the block-tridiagonal symmetric matrix*

$$H_{m,n} = \begin{pmatrix} \hat{A}_{m+1} & \hat{B}_{m+1} & & & \\ \hat{B}_{m+1}^\top & \hat{A}_{m+2} & \hat{B}_{m+2} & & \\ & \ddots & \ddots & \ddots & \\ & & \hat{B}_{n-3}^\top & \hat{A}_{n-2} & \hat{B}_{n-2} \\ & & & \hat{B}_{n-2}^\top & \hat{A}_{n-1} \end{pmatrix},$$

where the matrices $\hat{A}_i$ and $\hat{B}_i$ are given by

$$\hat{A}_i = D_{i-1} B_{i-1}^{-1} + B_i^{-1} A_i\ ,\ \hat{B}_i = -B_i^{-1}.$$

Connecting orbits with positive definite Hessian matrix will be interesting in the sequel. All of their points must be monotone, otherwise the Hessian is degenerate.

3.3. Minimizing orbits

Definition: We say that a (m,n)-segment of orbit is *minimizing* iff it is monotone and the corresponding matrix $H_{m,n}$ is definite positive. Then, we say that an orbit is minimizing iff it is monotone and all its Hessian matrices $H_{m,n}$ are positive definite.

Remarks:

i) So then, minimizing means non-degenerate minimum.

ii) As the eigenvalues of a matrix depend continuously on its components, if we have a minimizing segment of orbit, then another segment of orbit close enough to the first will also be minimizing.

iii) All the subsegments of a minimizing segment are also minimizing.

iv) A minimizing orbit for F is also a minimizing orbit for a power of F, because in the second case the chains are defined with more constraints. This is due to the fact that the primitive function of a power of F is the sum of the values of the primitive function on the points of the corresponding segment. ◁

3.3.1. The MMS method. In [21], MacKay, Meiss and Stark wrote a Hessian matrix using the Lagrangian generating function, but if we use 2.4 we see that the two matrices coincide, in the case that such a function exists. Then, although the Lagrangian generating function does not exist, we can define a (local) extremal behaviour of the orbits.

On the other hand, they describe a method to block-diagonalize that matrix. They found that the diagonal blocks associated to $H_{0,n}$ are given by the recurrence

$$\begin{cases} \hat{D}_1 = \hat{A}_1, \\ \hat{D}_i = \hat{A}_i - \hat{B}_{i-1}^\top \hat{D}_{i-1}^{-1} \hat{B}_{i-1} \quad (i = 2 \div n-1), \end{cases}$$

provided $\hat{D}_{i-1}$ is invertible. Then, the matrix $H_{0,n}$ is positive definite iff all the symmetric matrices $\hat{D}_i$ are positive definite.

3.3.2. 1 degree of freedom. If $d = 1$, then we can obtain a recurrence for the characteristic polynomials of $H_{0,i}$ $(i > 1)$: $p_{i-1}(x) = \det(xI - H_{0,i})$. Then:

$$\begin{cases} p_0(x) = 1, p_1(x) = x - \hat{a}_1; \\ p_i(x) = (x - \hat{a}_i) p_{i-1}(x) - \hat{b}_{i-1}^2 p_{i-2}(x) \quad (i > 2), \end{cases}$$

where the lower-case letters represent the corresponding 1×1 matrices. The sequence of polynomials $\{p_i, p_{i-1}, \ldots, p_1, p_0\}$ is a Sturm sequence for the polynomial p_i. In particular, all the eigenvalues of $H_{0,i}$ are different (and real, of course) and we can compute the number of positive eigenvalues:

> If $p_{i-1}(0) \neq 0$, the number of positive eigenvalues of $H_{0,i}$ is equal to the number of changes of sign in the sequence $\{p_0(0), p_1(0), \ldots, p_{i-1}(0)\}$.

This is a specific case of Sturm's theorem. Hence, we have to compute the number of changes of sign in the sequence

$$\begin{cases} r_0 = 1,\ r_1 = -\hat{a}_1; \\ r_i = -\hat{a}_i r_{i-1} - \hat{b}_{i-1}^2 r_{i-2} \quad (i > 1). \end{cases}$$

This fits into the MMS method, by defining $\forall i \geq 1\ \hat{d}_i = -\dfrac{r_i}{r_{i-1}}$.

3.4. Invariance of the extremal character of an orbit

We consider whether the extremal character of an orbit is independent of the variables in which we write our symplectomorphism. In fact, we shall see that it does not change under lifts and vertical translations, but it can change by conjugating with other symplectomorphisms. This is due to the concomitant distinction between x and y variables. For the sake of simplicity, we shall work in the monotone case. Otherwise, the orbits are degenerate as critical chains of the action.

Proposition 3 *Given a symplectomorphism $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$, the extremal character of monotone orbits does not change by (symplectic) vertical translations and lifts.*

Proof: We shall prove the first point, the second being similar (see [13]). So then, let $G = \tau_{\nabla l}$ be the vertical translation induced by the function $l : \mathbb{R}^d \to \mathbb{R}$, which defines a change of variables

$$\begin{cases} x = \bar{x}, \\ y = \bar{y} + \nabla l(\bar{x}). \end{cases}$$

Our symplectomorphism F written in the new variables if $\bar{F} = G^{-1} \circ F \circ G$, and it is given by

$$\begin{cases} \bar{x}' = f(\bar{x}, \bar{y} + \nabla l(\bar{x})), \\ \bar{y}' = g(\bar{x}, \bar{y} + \nabla l(\bar{x})) - \nabla l(f(\bar{x}, \bar{y} + \nabla l(\bar{x}))). \end{cases}$$

Recall that if the primitive function of F is S, then the primitive function of $\bar{F}$ is $S \circ G + l \circ q - l \circ f \circ G$, where q is the projection on the x-variables.

We consider now two corresponding orbits of F and $\bar{F}$. Hence, let (x_0, y_0) the initial point of an F-orbit and $(\bar{x}_0, \bar{y}_0) = G^{-1}(x_0, y_0)$ the initial point of the corresponding $\bar{F}$-orbit. We know that the extremal character of an F-orbit is given by the recurrence

$$\begin{cases} \hat{D}_1 = \hat{A}_1, \\ \hat{D}_i = \hat{A}_i - \hat{B}_{i-1}^{\top} \hat{D}_{i-1}^{-1} \hat{B}_{i-1} \quad (i \geq 2), \end{cases}$$

where $\hat{A}_i = D_{i-1} B_{i-1}^{-1} + B_i^{-1} A_i$ and $\hat{B}_i = -B_i^{-1}$. We should write the same sequence for an $\bar{F}$-orbit putting bars in all positions, but we shall write $\check{A}_i$ rather than $\hat{\bar{A}}_i$, etc. We

have to relate both sequences and see that the indexes of the matrices $\hat{D}_i$ and $\check{D}_i$ are the same. First of all, we relate $\mathrm{D}\bar{F}(\bar{z})$ and $\mathrm{D}F(z)$ by means of

$$\begin{pmatrix} \bar{A} & \bar{B} \\ \bar{C} & \bar{D} \end{pmatrix} = \begin{pmatrix} A + BL_0 & B \\ \bar{C} & -L_1 B + D \end{pmatrix},$$

where $L_0 = \mathrm{D}^2 l(\bar{x})$, $L_1 = \mathrm{D}^2 l(f(\bar{x}, \bar{y}))$, etc. Notice that monotonicity does not change under vertical translations.

Finally, we obtain that $\check{B}_i = \hat{B}_i$ and

$$\check{A}_i = (-L_i B_{i-1} + D_{i-1}) B_{i-1}^{-1} + B_i^{-1}(A_i + B_i L_i) = D_{i-1} B_{i-1}^{-1} + B_i^{-1} A_i = \hat{A}_i,$$

and the corresponding sequences of diagonal blocks $\hat{D}_i$ and $\check{D}_i$ coincide. □

Remark: A physical interpretation of this result is that the laws of *discrete mechanics* are independent of the coordinates in our configuration space and certain privileged observers. This is geometrically connected with the choice of a certain 1-form $\boldsymbol{\alpha} = y\ \mathrm{d}x$ in our phase space, and the distinction between x and y coordinates that it produces.◁

4. Invariant Lagrangian graphs

The first step in order to understand the properties of invariant Lagrangian manifolds is to study the simpler ones: the *invariant Lagrangian graphs* (in short, *i.L.g.*).

This section seeks to extend results reported by Mather [23], Herman [15] and MacKay, Meiss and Stark [21], obtained when using a (global) Lagrangian generating function. To a certain extent our results are more local, because we do not use the existence of this global function, and this allows us to study different regions in our phase space where some positivity conditions are satisfied.

4.1. Characterization of i.L.g.

Definition: Given a function $l : \mathbb{R}^d \to \mathbb{R}$, it is well known that the immersion

$$\begin{array}{rccc} \nu: & \mathbb{R}^d & \longrightarrow & \mathbb{R}^d \times \mathbb{R}^d \\ & x & \longrightarrow & (x, \nabla l(x)) \end{array}$$

defines a Lagrangian embedding of $\mathbb{R}^d$ into $\mathbb{R}^d \times \mathbb{R}^d$. The function l is known as the *generating function* of the *Lagrangian graph* $\mathcal{L}_{\nabla l} = \nu(\mathbb{R}^d) = \{y = \nabla l(x)\}$.

Definition: Given a symplectomorphism $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$, with $\mathrm{pf}(F) = S$, we can associate with the Lagrangian graph $\mathcal{L}_{\nabla l}$ a *critical function* $\hat{\Phi} : \mathbb{R}^{2d} \to \mathbb{R}$ by

$$\hat{\Phi}(x, y) = S(x, y) - (l(f(x, y)) - l(x)).$$

Its value on the graph is denoted by $\Phi(x) = \hat{\Phi}(x, \nabla l(x))$.

We can thus obtain the next proposition of characterization of invariant Lagrangian graphs.

Proposition 4 *Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism with* $\mathrm{pf}(F) = S$ *and $\mathcal{L} = \mathcal{L}_{\nabla l}$ be the Lagrangian graph given by $l : \mathbb{R}^d \to \mathbb{R}$. We define the functions Φ and $\hat{\Phi}$ as above. Then:*

(i) $\mathcal{L}$ is F-invariant $\Rightarrow$ Φ is constant
(Conserved quantity on an i.L.g.).

(ii) $\mathcal{L}$ is F-invariant $\Leftrightarrow$ $\forall x \in \mathbb{R}^d$ $\mathrm{D}\hat{\Phi}(x, \nabla l(x)) = 0$
(Characterization of i.L.g.).

Proof: The *invariance condition* is $\forall x \in \mathbb{R}^d \quad g(x, \nabla l(x)) = \nabla l(f(x, \nabla l(x)))$. The derivatives of $\hat{\Phi}$ are:

$$\frac{\partial \hat{\Phi}}{\partial x}(x, y) = \left(g(x, y)^\top - \frac{\partial l}{\partial x}(f(x, y)) \right) \cdot \frac{\partial f}{\partial x}(x, y) - y^\top + \frac{\partial l}{\partial x}(x),$$

$$\frac{\partial \hat{\Phi}}{\partial y}(x, y) = \left(g(x, y)^\top - \frac{\partial l}{\partial x}(f(x, y)) \right) \cdot \frac{\partial f}{\partial y}(x, y).$$

So, if $\mathcal{L}$ is F-invariant the two derivatives vanish on the graph and, in particular, the function Φ is constant:

$$\frac{\partial \Phi}{\partial x}(x) = \frac{\partial \hat{\Phi}}{\partial x}(x, \nabla l(x)) + \frac{\partial \hat{\Phi}}{\partial y}(x, \nabla l(x)) \cdot \frac{\partial^2 l}{\partial x^2}(x) = 0.$$

Conversely, if the derivatives vanish at a point $(x, \nabla l(x))$, then we obtain

$$g(x, \nabla l(x)) = \nabla l(f(x, \nabla l(x))),$$

because the rank of the matrix $(A(x, y) \;\; B(x, y))$ is maximal at all points. □

Remark: The first point can easily be generalized to any invariant exact Lagrangian manifolds of any exact symplectomorphisms. In fact, the whole proposition could also be generalized to any invariant exact Lagrangian manifolds by means of Weinstein's theorems [30], which enable us to send a tubular neighborhood of a Lagrangian manifold to a tubular neighborhood of the zero section of its cotangent bundle. For more information about this subject see [13]. ◁

4.2. Extremal character of an i.L.g.

Continuing with the above notation, a point of an i.L.g. $\mathcal{L} = \mathcal{L}_{\nabla l}$ is a fibered critical point of the critical function $\hat{\Phi}$, that is to say, for each point $x \in \mathbb{R}^d$

$$\frac{\partial \hat{\Phi}}{\partial y}(x, \nabla l(x)) = 0.$$

(The converse is true if F is monotone).

Definition: If all the points have the same character as critical points of the 'fibered' function, and then all the corresponding Hessian matrices have non-zero eigenvalues, we shall say that our graph is *non-degenerate*. In such a case, if all these matrices are positive definite we shall say that our graph is *minimizing*, and if all of them are negative definite we shall say that it is *maximizing*. Otherwise we shall say that it is *indefinite*.

4.3. Minimizing i.L.g.

Hence, the extremal character of the graph on each point $(x, \nabla l(x))$ is given by

$$\frac{\partial^2 \hat{\Phi}}{\partial y^2}(x, \nabla l(x)) = (D^\top(x) - B(x)^\top \, \mathrm{D}^2 l(\phi(x)))B(x),$$

where we write $\phi(x) = f(x, \nabla l(x))$, $A(x) = A(x, \nabla l(x))$, etc. If the graph is monotone, that is, it is included within a monotone region, the signature of such a matrix does not change if we multiply it by B^{-1} and $B^{-\top}$:

$$B(x)^{-\top} \frac{\partial^2 \hat{\Phi}}{\partial y^2}(x, \nabla l(x)) B(x)^{-1} = D(x)B(x)^{-1} - \mathrm{D}^2 l(\phi(x)).$$

By deriving the equalities $\phi(x) = f(x, \nabla l(x))$ and $\nabla l(\phi(x)) = g(x, \nabla l(x))$ we obtain

$$\begin{aligned} \mathrm{D}\phi \qquad &= A + B \, \mathrm{D}^2 l, \\ \mathrm{D}^2 l \circ \phi \, \mathrm{D}\phi &= C + D \, \mathrm{D}^2 l. \end{aligned}$$

Hence, as $D^\top A - B^\top C = I_d$, $D^\top B = B^\top D$ and $\mathrm{D}^2 l$ is symmetric, we obtain

$$\begin{aligned} B^{-1}A + \mathrm{D}^2 l \quad &= B^{-1}\mathrm{D}\phi, \\ DB^{-1} - \mathrm{D}^2 l \circ \phi &= (\mathrm{D}\phi)^{-\top} B^{-1}. \end{aligned}$$

Finally, we define the maps $E_1, E_2 : \mathbb{R}^d \to M_d(\mathbb{R})$ by $E_1(x) = B(x)^{-1}\mathrm{D}\phi(x)$ and $E_2(x) = (\mathrm{D}\phi(x))^{-\top} B(x)^{-1}$. They are symmetric and non-singular matrices, and they are related by

$$E_2(x) = B(x)^{-\top} E_1(x)^{-1} B(x)^{-1}.$$

Hence, if one is positive definite, the other one is also positive definite. Then, we obtain the following lemma (cf. [15, 16]).

Lemma 1 *Let $F : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism and $\mathcal{L} = \mathcal{L}_{\nabla l}$ be a monotone F-invariant Lagrangian graph. Let $E_1, E_2 : \mathbb{R}^d \to M_d(\mathbb{R})$ be the maps defined as above. Then:*

$$\mathcal{L} \text{ is minimizing} \Leftrightarrow \forall x \in \mathbb{R}^d \; E_1(x) \succ 0 \Leftrightarrow \forall x \in \mathbb{R}^d \; E_2(x) \succ 0.$$

4.4. Invariance of the extremal character of an i.L.g.

As we shall see, the extremal character of our graph does not change under vertical translations and lifts. Actually, we shall not see all of this, but sufficient for our purposes.

Proposition 5 *Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism with $\mathrm{pf}(F) = S$ and $\mathcal{L} = \mathcal{L}_{\nabla l}$ be an F-invariant Lagrangian graph generated by $l : \mathbb{R}^d \to \mathbb{R}$. Hence, the extremal character of the graph does not change after the following two steps of normal form:*

(i) projection of the zero-section,

(ii) simplification of the dynamics on that zero-section, via conjugation by a lift.

Proof: We shall prove the first step, the second being similar (see [13]). So then, the extremal character of our graph is given by the symmetric matrices

$$(D^\top(x) - B(x)^\top \, \mathrm{D}^2 l(\phi(x)))B(x).$$

If we perform a change of variables by means of the vertical translation

$$\begin{cases} x = \bar{x}, \\ y = \bar{y} + \nabla l(\bar{x}), \end{cases}$$

then in the new variables $(\bar{x}, \bar{y})$ the zero-section $\{\bar{y} = 0\}$ is fixed. After this projection, the character of the graph (the zero-section) is given by

$$\bar{D}(\bar{x})^\top \bar{B}(\bar{x}) = (D(x) - L_1(\bar{x}, 0)B(x))^\top B(x) = D(x)^\top B(x) - B(x)^\top \mathrm{D}^2 l(\phi(x))B(x),$$

and the character does not change (see the notation in 3.4). □

4.5. Orbits on minimizing invariant Lagrangian graphs

In this section we shall see that the orbits on a minimizing invariant Lagrangian graph are minimizing. This is the key result of the paper.

Theorem 1 *Let $F = (f, g) : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$ be a symplectomorphism and $\mathcal{L} = \mathcal{L}_{\nabla l}$ be a F-invariant Lagrangian graph given by $l : \mathbb{R}^d \to \mathbb{R}$. Then:*

If $\mathcal{L}$ is minimizing, then all the orbits on $\mathcal{L}$ are minimizing.

Proof: Before carrying out a complete proof we shall see what the key is to our constructions. So, let S be the primitive function of F, $\hat{\Phi}$ be the function associated to the graph, and $C \in \mathbb{R}$ be the corresponding conserved quantity: $\forall x \in \mathbb{R}^d \ \hat{\Phi}(x, \nabla l(x)) = C$. As $\mathcal{L}$ is minimizing, then in a tubular neighborhood of the graph we have $\hat{\Phi}(x, y) \geq \hat{\Phi}(x, \nabla l(x)) = C$. Let $(\mathbf{x}, \nabla l(\mathbf{x}))$ be any point on $\mathcal{L}$. Fix $n > 1$, let $\mathcal{O} = ((\mathbf{x}_i, \mathbf{y}_i) = F^i(\mathbf{x}, \nabla l(\mathbf{x})))_{i=0 \div n-1}$ be a segment of orbit and $\mathcal{C} = (x_i, y_i)_{i=0 \div n-1}$ be any F-chain from $\mathbf{x}_0$ to $\mathbf{x}_n = f(\mathbf{x}_{n-1}, \mathbf{y}_{n-1})$, close enough to the graph. So then:

$$\begin{aligned}
\boldsymbol{S}_{0,n}(\mathcal{C}) &= \sum_{i=0}^{n-1} S(x_i, y_i) \\
&\overset{\text{chain}}{=} \sum_{i=0}^{n-1} (S(x_i, y_i) - l(f(x_i, y_i)) + l(x_i)) \ + l(\mathbf{x}_n) - l(\mathbf{x}_0) \\
&= \sum_{i=0}^{n-1} \hat{\Phi}(x_i, y_i) \ + l(\mathbf{x}_n) - l(\mathbf{x}_0) \\
&\geq nC + l(\mathbf{x}_n) - l(\mathbf{x}_0) \\
&= \sum_{i=0}^{n-1} \hat{\Phi}(\mathbf{x}_i, \mathbf{y}_i) \ + l(\mathbf{x}_n) - l(\mathbf{x}_0) \\
&= \boldsymbol{S}_{0,n}(\mathcal{O}).
\end{aligned}$$

Hence, our connecting orbit minimizes the action on the chains. Notice that the restriction of the action to the set of F-chains is fundamental.

We have to improve this result and show that any segment of orbit is, in fact, a non-degenerate minimum of the corresponding action. We do not need global conditions. Thanks to the invariance of the extremal character of orbits (3.4) and graphs (4.4) under vertical translations, we can restrict our attention to the case in which our graph is the zero-section. In such a case, our symplectomorphism is given by

$$\begin{cases} x' = \phi(x) + B(x)y + \dots, \\ y' = \mathrm{D}\phi(x)^{-\top} y + \dots, \end{cases}$$

where '...' means terms in upper orders in y. Hence we have

$$\mathrm{D}F(x,0) = \begin{pmatrix} A(x) & B(x) \\ 0 & D(x) \end{pmatrix},$$

where $A(x) = \mathrm{D}\phi(x)$, $D(x) = A(x)^{-\top}$, and $A(x)B(x)^{\top} = B(x)A(x)^{\top}$ (2.1). Therefore

$$\frac{\partial^2 \hat{\Phi}}{\partial y^2}(x,0) = \frac{\partial^2 S}{\partial y^2}(x,0) = D(x)^{\top} B(x) = A(x)^{-1} B(x).$$

Hence, our graph is minimizing iff $\forall x \in \mathbb{R}^d$ $A(x)^{-1}B(x) \succ 0$, where $\succ$ is the Loewner partial order (and $S \succ 0$ means that the symmetric matrix S is positive definite). In fact, as the symmetric matrices $A^{-1}B = B^{\top}A^{-\top}$ and $B^{-\top}A^{-1} = A^{-\top}B^{-1}$ have the same inertia, we have

$$A^{-1}B \succ 0 \Leftrightarrow B^{-1}A \succ 0 \Leftrightarrow A^{-\top}B^{-1} \succ 0 \Leftrightarrow AB^{\top} \succ 0,$$

applying that $M \succ 0 \Rightarrow M^{-1} \succ 0$, for any symmetric matrix M.

Now, let $(x, \nabla l(x))$ be any point on the graph. We shall write $A_i = A(\phi^i(x))$, $B_i = B(\phi^i(x))$ and $D_i = D(\phi^i(x)) = A_i^{-\top}$. We shall use the MMS method (3.3.1). In this case we have

$$\hat{A}_i = A_{i-1}^{-\top} B_{i-1}^{-1} + B_i^{-1} A_i \succ 0.$$

We shall prove by induction that $\hat{D}_i = B_i^{-1}A_i + K_i$, where $K_i \succ 0$, for any $i \geq 1$.

- For $i = 1$ we have $\hat{D}_1 = B_1^{-1}A_1 + K_1$, where $K_1 = A_0^{-\top}B_0^{-1} \succ 0$, and the property holds.
- Assume that the property holds for $i-1$. So then

  $$\hat{D}_{i-1} = B_{i-1}^{-1}A_{i-1} + K_{i-1} \succ B_{i-1}^{-1}A_{i-1} \succ 0$$

  and

  $$B_{i-1}\hat{D}_{i-1}B_{i-1}^{\top} \succ A_{i-1}B_{i-1}^{\top} \succ 0.$$

  On the other hand, we have $\hat{D}_i = B_i^{-1}A_i + K_i$, where

  $$K_i = A_{i-1}^{-\top}B_{i-1}^{-1} - B_{i-1}^{-\top}\hat{D}_{i-1}^{-1}B_{i-1}^{-1} = A_{i-1}^{-\top}B_{i-1}^{-1} - (B_{i-1}\hat{D}_{i-1}B_{i-1}^{\top})^{-1}.$$

  Finally,

  $$K_i \succ A_{i-1}^{-\top}B_{i-1}^{-1} - B_{i-1}^{-\top}A_{i-1}^{-1} = 0.$$

  In summary, all the matrices $\hat{D}_i$ are positive definite. □

5. Converse KAM theory

Some non-existence criteria of invariant tori are founded on some type of positivity. We have followed [15] rather than [21], that is, we have studied monotone positive cases rather than twist cases.

5.1. Set up

We shall work on the annulus, $\mathbb{A}^d$. Let $F : \mathbb{A}^d \to \mathbb{A}^d$ be a symplectomorphism. We shall consider its lift $\tilde{F} : \mathbb{R}^{2d} \to \mathbb{R}^{2d}$, which is a symplectomorphism with primitive function $S : \mathbb{R}^{2d} \to \mathbb{R}$.

Let $\mathcal{T}$ be an F-invariant Lagrangian horizontal torus, that is, an i.L.g.

$$y = a + \nabla \bar{l}(x)$$

where $a \in \mathbb{R}^d$ and $l : \mathbb{T}^d \to \mathbb{R}$. So, the generating function (on $\mathbb{R}^d$) of the graph is $l(x) = a \cdot x + \bar{l}(x)$. That is, $\mathcal{T} = \mathcal{L}_{\nabla l}$.

Let $\phi : \mathbb{T}^d \to \mathbb{T}^d$ be the dynamics on the torus, that is to say, the *diffeomorphism* given by $\phi(x) = f(x, \nabla l(x))$. We shall also write $A(x) = A(x, \nabla l(x)), B(x) = B(x, \nabla l(x)$, etc. Suppose also that $\mathcal{T}$ is monotone ($\forall x \in \mathbb{T}^d \ |B(x)| \neq 0$).

Remark: In the course of this paper we have supposed that our invariant Lagrangian tori are graphs. Unfortunately, there is no equivalent to higher dimensions of the next Birkhoff's theorem, for $d = 1$:

> Let $F : \mathbb{A} \to \mathbb{A}$ be a C^1 monotone symplectomorphism, satisfying
>
> $$\sup_{z \in \mathbb{A}} (|B^{-1}(z)A(z)|, |D(z)B^{-1}(z)|) < \infty.$$
>
> Then, any C^0 F-invariant torus homotopic to the circle $\{y = 0\}$, called *invariant rotational curve* (in short, *i.r.c.*), is the graph $\mathcal{L}_\psi$ of a certain continuous function $\psi \in C^0(\mathbb{T}^1, \mathbb{R})$.

For the proof of this theorem see [8, 14, 24]. Herman [15] has perturbative generalizations to higher dimension of this theorem. See [7] for a generalization of this 'second' Birkhoff's theorem to optical Hamiltonian diffeomorphisms. ◁

5.2. Monotone positivity implies minimizing

The following theorem completes the one above, and gives a sufficient condition for a minimizing i.L.g. (cf. [15, 16]).

Theorem 2 *Let $F : \mathbb{A}^d \to \mathbb{A}^d$ be a symplectomorphism, and $\mathcal{T} = \mathcal{L}_{\nabla l}$ be an F-invariant Lagrangian horizontal torus given by $l(x) = a \cdot x + \bar{l}(x)$ (using the notation above). Then:*

> *If $\mathcal{T}$ is monotone positive, then $\mathcal{T}$ is minimizing.*

Proof: Suppose that the graph is monotone ($+_a$), that is, $\forall x \in \mathbb{T}^d$ $B^{-1}(x)A(x) \succ 0$ (2.3). Let x_0 be the minimum of the periodic function $\bar{l}$. Then:

$$\mathrm{D}^2 l(x_0) = \mathrm{D}^2 \bar{l}(x_0) \succeq 0.$$

Hence:

$$E_1(x_0) = B^{-1}(x_0)A(x_0) + \mathrm{D}^2 l(x_0) \succ 0.$$

Finally, since E_1 is non-singular everywhere, it is also definite positive everywhere, and the graph is minimizing.

If we suppose that the graph is monotone ($+_d$) the proof is similar. We have to take the preimage by ϕ of the maximum of $\bar{l}$ to prove that E_2 is positive definite everywhere. □

Remarks:

i) We have that $\forall x \in \mathbb{T}^d$

$$E_1(x) + E_2(\phi^{-1}(x)) = D(\phi^{-1}(x))B^{-1}(\phi^{-1}(x)) + B^{-1}(x)A(x) = \hat{A}(x).$$

Hence, if our graph is minimizing (for instance, if our graph is monotone positive) then these matrices are all positive definite. These matrices appear on the diagonal of the second derivative of the action, and this result was previously obtained. In particular, if $\hat{A}(x,y) \preceq 0$ then there is no minimizing invariant graph through (x,y). For $d = 1$, this coincides with the first step in the *Lipschitz criterion* for non-existence of invariant graphs [20, 25].

ii) Herman [15, 16] obtained an upper bound for the slope of the monotone positive i.L.g.. $y = \psi(x) = \nabla l(x)$. From the inequality

$$-B^{-1}(x)\ A(x) \prec \mathrm{D}\psi(x) \prec D(\phi^{-1}(x))\ B^{-1}(\phi^{-1}(x)),$$

which is satisfied $\forall x \in \mathbb{T}^d$, he obtained

$$\|\rho(\mathrm{D}\psi)\|_\infty \leq \max(\|\rho(B^{-1}A)\|_\infty, \|\rho(DB^{-1})\|_\infty)$$

where $\|\cdot\|_\infty$ means the sup-norm of a function defined on $\mathbb{T}^d$ and ρ means the spectral radius of a matrix. This is the main result of BHM theory, and we have used its proof in relating it with Converse KAM theory.

This upper bound is a little stronger than the previous one, because we have a stronger hypothesis. Herman also proved that if our exact symplectomorphism is $\mathcal{C}^1$ and monotone globally positive and the invariant torus is $\mathcal{C}^0$-Lagrangian (see [15, 16] for definitions and proofs). It is a generalization to a higher dimension of a Birkhoff's theorem, for $d = 1$ [8], that gives us bounds of the slope of i.r.c..

In the case $d = 1$, such bounds give Lipschitz cones, and inside them lies the i.r.c.. This is the core of the *cone-crossing* criterion for the non-existence of i.r.c. performed by MacKay and Percival [20], and first used by Herman [14] and Mather [24]. On the other hand, Newman and Percival [27] and, independently, Aubry and co-workers [4, 5], used criteria connected with action principles. The equivalence of both methods was proved in [20].

iii) A similar proposition was also proved by Herman [15, 16] and by MacKay, Meiss and Stark [21] using different assumptions. They need the generating function (and impose twist conditions on the symplectomorphism). We believe that many results can be proved without using a global generating function which satisfies some strong conditions of positivity.

iv) Notice that E_1 is $B^{-1}A$ and E_2 is DB^{-1} *after projection* of the graph on the zero-section. So then, the graph is minimizing iff when we project it on the zero-section then it is monotone positive in the two senses. ◁

5.3. Non-existence criterion

In [21], a *variational criterion* for the non-existence of invariant Lagrangian graphs was derived. We can write it as follows. We shall use the same notation as in the proposition.

> *If the orbit by z yields on a monotone positive region and it has a non-minimizing segment, then it does not lie on any invariant Lagrangian horizontal torus included within such a region.*

In order to determine whether a segment is minimizing we shall use the MMS method (3.3.1). Of course, if we take segments of length 1 we obtain rather crude estimates.

In [21], the test was applied to the Froeschlé map [11], which is a 4D twist map (given by a Lagrangian generating function). However, we have seen that we should not be so restrictive, and we can apply their methods to other examples.

6. Examples

6.1. Generalized standard-like maps

A *generalized standard-like map* [19] is a symplectomorphism in $\mathbb{T}^d \times \mathbb{R}^d$ given by two *potentials*, $V : \mathbb{T}^d \to \mathbb{R}$ and $W : \mathbb{R}^d \to \mathbb{R}$, by means of

$$\begin{cases} y' = y - \nabla V(x), \\ x' = x + \nabla W(y') \pmod 1. \end{cases}$$

Its inverse is given by

$$\begin{cases} y = y' + \nabla V(x), \\ x = x' - \nabla W(y') \pmod 1. \end{cases}$$

Its primitive function is

$$S(x, y) = (y - \nabla V(x)) \cdot \nabla W(y - \nabla V(x)) - W(y - \nabla V(x)) - V(x)$$

(notice that it is 1-periodic in all its x-variables). The Jacobian matrix is

$$\mathrm{D}F(x, y) = \begin{pmatrix} I_d - \mathrm{D}^2 W(y')\, \mathrm{D}^2 V(x) & \mathrm{D}^2 W(y') \\ -\mathrm{D}^2 V(x) & I_d \end{pmatrix}.$$

Moreover, $\hat{A}(x, y) = \mathrm{D}^2 W(y)^{-1} + \mathrm{D}^2 W(y')^{-1} - \mathrm{D}^2 V(x)$ and $\hat{B}(x, y) = -\mathrm{D}^2 W(y')^{-1}$ (when this is possible).

Examples:

1) If we choose $V(x) = 0$ we obtain an integrable map, and all the tori $\{y = y_0\}$ are invariant. The corresponding frequency vectors are $\omega(y_0) = \nabla W(y_0)$. The extremal character of the orbits depends on $\mathrm{D}^2 W(y_0)$. If $\mathrm{D}^2 W(y_0) \succ 0$ the orbits are minimizing and if $\mathrm{D}^2 W(y_0) \prec 0$ the orbits are maximizing. Indefinite orbits appear when $\mathrm{D}^2 W(y_0)$ is indefinite.

2) If we take $W(y) = \frac{1}{2} y^2$ we obtain a *standard-like map*, which is generated by the Lagrangian generating function

$$L(x, x') = \frac{1}{2}(x' - x)^2 - V(x)$$

and it is given by

$$\begin{cases} y' = y - \nabla V(x), \\ x' = x + y - \nabla V(x) \pmod 1. \end{cases}$$

The standard-like maps are monotone ($+_d$), and twist. They are discrete models of the Newton's second law, because

$$x'' - 2x' + x = -\nabla V(x').$$

Hence, an orbit is determined by its sequence of angles. ◁

If the potential W is a strictly convex function ($\mathrm{D}^2 W(y) \succ 0$ for any point $y \in \mathbb{R}^d$) then our symplectomorphism is monotone ($+_d$). If this is not the case, we can study the monotone positive region:

$$\mathcal{M}_+ = \{(x, y) \in \mathbb{T}^d \times \mathbb{R}^d \mid \mathrm{D}^2 W(y - \nabla V(x)) \succ 0\}.$$

Of course, we can do the same with the monotone negative region $\mathcal{M}_-$.

Applying the first step in the variational criterion, if there is an i.L.g. inside $\mathcal{M}_+$ through a point $(x, y) \in \mathcal{M}_+$, then the corresponding orbit must be minimizing and, in particular, the matrix $\hat{A}(x, y)$ must be positive definite. As an i.L.g. intersects each fiber $\{x = x_0\}$, we can focus on one in particular. For instance, we can take x_0 as a critical point of V, and then $\hat{A}(x_0, y) = 2\,\mathrm{D}^2 W(y)^{-1} - \mathrm{D}^2 V(x_0)$.

Of course, stronger results may be obtained by iterating, with the aid of a computer. For instance, suppose our map be positive definite. Then, we can throw out the pieces of the phase space where the points are not minimizing after a finite number of iterations. We are sure that in these pieces there are no invariant tori. On the other hand, minimizing orbits not only correspond to invariant tori, but also to minimizing periodic orbits, cantori, etc. This is the philosophy in [21]. To obtain rigorous results we should use techniques of interval analysis [20].

We shall apply the method to different generalized standard maps.

6.2. Some 2D examples

We shall consider $d = 1$, that is, an *area preserving map* (in short, *a.p.m.*), and the potential V given by

$$V(x) = -\frac{K}{(2\pi)^2}\cos(2\pi x),$$

where K is a positive perturbative parameter. In these examples we shall apply:

- the first step in the variational criterion, in order to obtain rather crude estimates of the critical value of K in which all the i.r.c. have been broken;
- the MMS method to a region of the phase space, in order to check in which parts of the phase space invariant tori do not exist and in which parts their existence is possible.

We have taken different values of K to illustrate the minimizing and maximizing regions. We have taken the piece of the cylinder, $C_{[-1,1]} = \mathbb{T} \times [-1, 1]$, and we have compared:

- dynamics, taking 1024 points and iterating each of them 1024 times;
- extremal character of the orbits, applying the MMS method to segments of length 128 and drawing the initial points with a colour, which depends on the number of positive eigenvalues of the corresponding Hessian matrix (the scale appears in *figure 1*);
- minimizing and maximizing regions, by choosing the corresponding points from the previous picture.



Figure 1. Scale from minimizing to maximizing character. The middle of the scale represents minimaximizing character.

Examples:

1) *The standard map.* The *standard map* or *Taylor-Chirikov map* [10] has $W(y) = \frac{1}{2}y^2$ and it is monotone $(+_d)$. Then:

$$\hat{A}(x, y) = 2 - V''(x) = 2 - K\cos(2\pi x).$$

As $K > 0$, $A(x, y)$ takes his smaller value for $x = 0$, being $\hat{A}(0, y) = 2 - K$. So then, as in the first step in Mather's calculations [24]:

If $K \geq 2$, there are no i.L.g..

If we take a segment of length 2, then we must take into account $\hat{D}_1(0, y) = 2 - K$ and

$$\hat{D}_2(0, y) = 2 - K\cos(2\pi y) - \frac{1}{2 - K}.$$

If $0 < K < 2$ then $\hat{D}_1 > 0$ and there are no i.L.g. if $\hat{D}_2 \leq 0$. The maximum value of $\hat{D}_2$ is taken for $y = \frac{1}{2}$ and it is $2 + K - \frac{1}{2-K}$. Finally, we improve the previous bound and we find there are no i.L.g. if $K \geq \sqrt{3}$. We could improve the bounds by taking into account longer segments.

A better bound is obtained by improving the Lipschitz cone. For instance, Mather [24] obtained the bound $K \geq 4/3$ taking into account segments of length 2 (this kind of bound was generalized by Herman [15] to higher dimensions). Later, MacKay and Percival [20] refined it to obtain $K \geq 63/64$. This refinement is an example of a computer assisted proof. Finally, Jungreis [17] also performed a method for proving (computer assisted) that the standard map has no invariant circles for $K = 0.9718$. These bounds are in accordance with the result of Greene [12], who estimated (not rigorously) the bound $K > 0.971635406$, by means of the residue criterion.

Figure 2 shows how the invariant tori disappear when we increase the parameter K.

2) *The quadratic standard map.* We consider $W(y) = \frac{1}{3}y^3$. This is an example of non-monotone a.p.m., because the monotonicity condition fails on the curve $y = V'(x)$. Notice that the union of non-monotone orbits, that is, the set of images and preimages of $y = V'(x)$, has zero measure. As $K > 0$, the points over $y = V'(x)$ are monotone $(+_d)$, and the points below $y = V'(x)$ are monotone $(-_d)$. The a.p.m. has no Lagrangian generating function.

As

$$\hat{A}(0, y) = \frac{1}{y} - K, \ \hat{A}(\frac{1}{2}, y) = \frac{1}{y} + K$$

then

there is no monotone $(+_d)$ i.L.g. through any point $(0, y)$ with $y \geq \frac{1}{K}$, and
there is no monotone $(-_d)$ i.L.g. though any point $(\frac{1}{2}, y)$ with $y \leq -\frac{1}{K}$.

Notice in *figure 3* that the non-monotone i.r.c. are more robust that their monotone counterparts, which seem to be graphs. Moreover, the non-monotone curves, which have folds, prevent the mixing between the monotone positive and monotone negative regions. When they break, mixing is possible. Then, we see that these curves are minimaximizing, in the sense that 'half' of the eigenvalues of the Hessian matrix are positive. Almost all i.r.c. are, in fact, monotone, if suitable coordinates are chosen (cf. [29]). We should adapt the coordinates to dynamics. ◁



Figure 2. Destruction of the i.r.c. in the standard map, when we increase the parameter K. The values of K are 0.9 and 1, respectively. The windows represent the cylinder $\{y \in [-1,1]\}$, where $x \in [0,1]$ is the angle variable. Recall that the last i.r.c. breaks for $K \simeq 0.971635406$.



Figure 3. Evolution of i.r.c for the quadratic standard map. The parameters are $K = 0.7$ and $K = 1.5$. The windows represent the cylinder $\{y \in [-1, 1]\}$, where $x \in [0, 1]$ is the angle variable. Non-monotone i.r.c. are stronger than monotone ones, and they are 'meandering'.

6.3. Around an elliptic fixed point

We can also apply these methods to the study of a neighborhood of an elliptic fixed point, by means of suitable changes of variables. For the sake of simplicity we shall consider the 2D case.

Suppose we have a symplectomorphism $F : \mathbb{R}^2 \to \mathbb{R}^2$, being the origin an elliptic fixed point. We consider the polar symplectic change of variables

$$\begin{array}{rccl} P: & \mathbb{S} \times \mathbb{R}^*_+ & \longrightarrow & \mathbb{R}^2 \\ & (\theta, I) & \longrightarrow & (x = \sqrt{2I}\cos\theta, y = \sqrt{2I}\sin\theta). \end{array}$$

In order to carry out the calculations, we have to consider the differential of $P^{-1} \circ F \circ P$. Note that it is not necessary to perform the change of variables and that it is sufficient to consider the matrices

$$M(x,y) = \begin{pmatrix} \frac{-\bar{y}}{\bar{x}^2+\bar{y}^2} & \frac{\bar{x}}{\bar{x}^2+\bar{y}^2} \\ \bar{x} & \bar{y} \end{pmatrix} \begin{pmatrix} a(x,y) & b(x,y) \\ c(x,y) & d(x,y) \end{pmatrix} \begin{pmatrix} -y & \frac{x}{x^2+y^2} \\ x & \frac{y}{x^2+y^2} \end{pmatrix},$$

being $(\bar{x}, \bar{y}) = F(x,y)$.

Example: In *figure 4*, we consider the standard map with $K = 1$. We also show dynamics, extremal character and minimizing regions, as in the previous subsection. The extremal character and the minimizing segments of orbits (of length 128) are computed in symplectic polar coordinates. Around the elliptic fixed point the map is monotone positive, except for the vertical axis. We have counted the number of non-negative eigenvalues of the Hessian matrices.



Figure 4. Around the origin, which is an elliptic fixed point of the standard map for $K = 1$. The window represents the square $[-0.5, 0.5] \times [-0.5, 0.5]$.

In these pictures, the resonance zone associated with the elliptic fixed point has been roughly bounded. The resonance zone, whose points surround the elliptic fixed point and do not escape, is bounded by an invariant curve. For a more accurate study of this 'last' invariant curve see the paper by Simó and Treschev [28]. ◁

6.4. A 4D example

Now, we consider $d = 2$ and the potential V given by

$$V(x_1, x_2) = -\frac{1}{(2\pi)^2}(K_1 \cos(2\pi x_1) + K_2 \cos(2\pi x_2) + \lambda \cos(2\pi(x_1 + x_2)).$$

We can consider different potentials $W(y_1, y_2) = W(y_1) + W(y_2)$, corresponding with the couplings (via λ) of two generalized standard maps. The Froeschlé map [11] corresponds to $W(y_1, y_2) = \frac{1}{2}(y_1^2 + y_2^2)$. For the next example and other similar ones see [13].

Example: We have considered a *standard×exponential Froeschlé map*, which is given by $W(y_1, y_2) = \frac{1}{2}y_1^2 + \exp(y_2)$. It is a monotone positive symplectomorphism, but it is not generated by a Lagrangian generating function. Since

$$\hat{A}(0, 0, y_1, y_2) = \begin{pmatrix} 2 - (K_1 + \lambda) & -\lambda \\ -\lambda & 2e^{-y_2} - (K_2 + \lambda) \end{pmatrix},$$

there are no i.L.g. if $K_1 + \lambda \geq 2$. Otherwise, there are no i.L.g. through any point $(0, 0, y_1, y_2)$ with

$$y_2 \geq \log\left(\frac{2(2 - (K_1 + \lambda))}{2(K_2 + \lambda) - K_1 K_2 - (K_1 + K_2)\lambda}\right).$$

In *figure 5* we show the extremal character of the points from a piece of the symmetry plane $\{x_1 = 0, x_2 = 0\}$ and we have chosen the minimizing ones.



Figure 5. Box $\{(y_1, y_2) \in [-1, 1] \times [-1, 1]\}$ in the slice $\{(x_1, x_2) = (0, 0)\}$, for the standard×exponential Froeschlé map. The parameters are $K_1 = 0.1$, $K_2 = 0.2$ and $\lambda = 0.1$. Notice that the number of invariant tori decreases when we increase y_2.

The channels in the second picture above are similar to those obtained in [21] for the Froeschlé map. They are related to the resonances associated with elliptic periodic orbits. ◁

Conclusions and perspectives

We have introduced variational principles for the orbits of any symplectomorphism defined on $\mathbb{R}^d \times \mathbb{R}^d$ (or, in general, for exact symplectomorphisms defined on the cotangent bundle of a manifold). The key point lies in using the primitive function and defining the action on the set of chains. Then, we have given a non-existence criteria of monotone positive invariant tori, because their orbits are minimizing. These methods also work for a broader class of maps: the symplectic skew-products [13].

In order to study the break-down of a certain invariant torus, we can apply criteria similar to Greene's [12], but instead of using the dynamic character of periodic orbits (given by their Floquet multipliers) we can use their extremal character.

For further information on these questions see [13].

Acknowledgements

My special thanks are due to C. Simó for his suggestions and encouragement. I would also like to thank R.S. MacKay for helpful discussions.

The research has been supported by DGICYT grant PB 94–0215 (Spain). Partial support from the EC grant ERBCHRXCT940460 and the Catalan grant CIRIT 1996S0GR-00105 is also acknowledged.

References

[1] Arnold V I 1963 *Russ. Math. Surveys* **18** 9
[2] Arnold V I 1976 *Les méthodes mathemátiques de la mécanique classique* (Moscow: Mir)
[3] Arnold V I, Kozlov V V and Neishtadt A I 1988 *Dynamical Systems III* (Springer-Verlag)
[4] Aubry S, Le Daeron P Y and André G 1982 *Classical ground-states of a 1-D model for incommensurate structures* (CEN Saclay)
[5] Aubry S 1983 *Physica D* **7** 240
[6] Avez A 1985 *Differential Calculus.* (New York: Wiley)
[7] Bialy M and Polterovich L 1992 *Math. Ann* **292** 619
[8] Birkhoff G D 1920 *Acta Mathematica* **43** 1 or *Collected Math. Papers* vol 2 p 111
[9] Broer H W, Huitema G B and Sevryuk M B *Quasi-periodic motions in families of dynamical systems. Order amidst chaos* (Lecture Notes in Math. **1645**)
[10] Chirikov B V 1979 *Phys. Rep.* **52** 265
[11] Froeschlé C 1972 *Astron. Astrophys.* **16** 172
[12] Greene J M 1979 *J. Math. Phys.* **20** 1183
[13] Haro A 1998 *The primitive function of an exact symplectomorphism* (Ph. D. Thesis, Universitat de Barcelona). `www.maia.ub.es/dsg/1998/index.html`.
[14] Herman M R 1983 *Sur les courbes invariantes per les différentiable par les difféomorphismes de l'anneau* vol 1 (Astérisque 103-104)
[15] Herman M R 1988 *École Polytechnique* **XIV** (séminaire 1987-1988)
[16] Herman M R 1989 *Publ. Math. I.H.E.S.* **70** 47
[17] Jungreis I 1991 *Ergodic Theory Dyn. Syst.* **11** 79
[18] Kolmogorov A N 1954 *Dokl. Akad. Nauk. SSR* **98** 469
[19] MacKay R S 1982 *Renormalization in area preserving maps* (Princeton thesis) (World Scientific 1993)

[20] MacKay R S and Percival I C 1985 *Commun. Math. Phys.* **98** 469
[21] MacKay R S, Meiss J D and Stark J 1989 *Nonlinearity* **2** 555
[22] Mather J N 1982 *Topology* **21** 457
[23] Mather J N 1984 *Letter to R.S. MacKay*
[24] Mather J N 1984 *Ergodic Theory Dyn. Syst.* **2** 301
[25] Meiss J D 1992 *Rev. Mod. Phys* **64** 795
[26] Moser J 1962 *Nach. Akad. Wiss. (Gottingen)* Math. Phys. Kl. II 1, 1
[27] Newman R A P C and Percival I C 1983 *Physica D* **6** 249
[28] Simó C and Treschev D V 1998 *Evolution of the 'last' invariant curve in a family of area preserving maps* Preprint
[29] Simó C 1998 *Invariant curves of perturbations of non twist integrable area preserving maps* Preprint
[30] Weinstein A 1977 *CBMS reg. conf. ser. Math.* **29** (Providence: AMS)